



---

# Official Grain Grading Guide

August 1, 2011

## Regional offices

Western Region .....Phone: 604-666-2028.....Fax: 604-666-8703  
Central Region.....Phone: 807-626-1401 .....Fax: 807-623-8701  
Eastern Region.....Phone: 514-283-7426.....Fax: 514-283-7699

## Service centers

Calgary .....Phone: 403-292-4211 .....Fax: 403-292-5075  
Saskatoon .....Phone: 306-975-5714 .....Fax: 306-975-4258  
Weyburn .....Phone: 306-848-3350.....Fax: 306-848-3353  
Winnipeg.....Phone: 204-983-2790.....Fax: 204-984-5131

**Quality • Service • Innovation**



---

## 25. Sample feed grain

---

### Determination of dockage

Samples are graded using procedures for sample feed grain when they exceed the Broken tolerances for Mixed Grain.

▲ **Important:** When a sample is to be graded as sample feed grain,

1. Return dockage to the cleaned sample.
2. Begin cleaning and dockage assessment using the procedures described in this section.

Dockage is assessed to the nearest 0.1% on all samples.

### Normal cleaning procedures

1. Using a Boerner-type divider, divide the uncleaned sample to obtain a representative portion.
  - Official samples should be at least 900 grams.
  - Unofficial samples must be at least 750 grams.
2. Sieve the sample over the No. 4.5 round-hole hand sieve, using approximately 250 g at a time.

### Composition of dockage

Dockage contains

- Material that passes through the No. 4.5 round-hole sieve
- Material removed by special cleaning for grade improvement

### Cleaning for grade improvement

If the grade of a sample can be improved by additional cleaning, perform the cleaning and add the additional material to dockage. Cleaning for grade improvement can be done at any time after the cleaning assessment has been completed.

1. After the cleaning assessment has been completed, use a No. 6 or a No. 5 buckwheat hand sieve to remove large seeds. Large seeds are removed if their total weight exceeds 3% of the cleaned sample by weight.
2. Record the additional cleaning and dockage on inspection records.

## Primary and export grade determinants tables

### Sample feed grain

Grade name	Foreign material												
	Ergot %	Excreta %	Flaxseed %	Fusarium %	Large material				Sclerotinia %	Soft earth pellets %	Stones %	Wild oats %	Total large material and wild oats %
					Wheat heads %	Knuckles and straw %	Large weed seeds %	Total %					
<b>Sample feed grain</b>	<b>0.33</b>	0.03	5	4	10	4	3	10	<b>0.25</b>	<b>0.33</b>	0.1	49	50
	<i>Sample Feed Grain, Ergoty</i>	<i>Sample (with components named)</i>	<i>Sample (with components named)</i>	<i>Sample (with components named)</i>	<i>Sample (with components named)</i>	<i>Sample (with components named)</i>	<i>Sample (with components named if not removable with approved sieves)</i>	<i>Sample (with components named)</i>	<i>Sample Feed Grain, Sclerotinia</i>	<i>Sample, Feed Grain, Soft Earth Pellets</i>	<i>2.5% or less— Sample Feed Grain, Stones Over 2.5%— Sample Feed Grain, Sample Salvage</i>	<i>Mixed Feed Oats</i>	<i>Sample (with components named)</i>

Large weed seeds includes other grains such as peas, corn, domestic buckwheat, etc.  
 Samples containing over 3.0% by weight are graded *Sample* and the components are named.  
 Soft earth pellets are included in total of large seeds.

Grade name	Damage		
	Broken grain %	Fireburnt %	Heated %
<b>Sample feed grain</b>	50	2	5
	<i>Sample Broken Grain (class on request)</i>	<i>Sample Feed Grain, Fireburnt</i>	<i>Sample Feed Grain, Heated</i>