

17. Corn

Determination of dockage	17-2
Definitions.....	17-2
Dockage not reported.....	17-2
Normal cleaning procedures:.....	17-2
Composition of dockage.....	17-3
Estimating test weight of well-matured corn after drying	17-4
Grading	17-5
Important definitions	17-5
Net weight of sample	17-5
Hazardous substances in samples	17-5
Processed sample.....	17-5
Rounding rules	17-5
Non-registered varieties	17-5
Representative portions for grading	17-6
Grading factors	17-7
Blue-eye mould (BEM).....	17-7
Caramelized kernels	17-7
Classes.....	17-7
Contaminated grain	17-7
Cracked corn and foreign material (CCFM)	17-8
Damage (DMG)	17-8
Earth pellets (EP)	17-8
Excreta (EXCR).....	17-8
Extraneous material	17-9
Fertilizer pellets (FERT PLTS)	17-9
Fireburnt (FBNT)	17-9
Foreign material (FM)	17-9
Heated (HTD).....	17-9
Odour (ODOR)	17-10
Rotted (ROT)	17-10
Stones (STNS).....	17-10
Test weight (TWT).....	17-12
Treated seed and other chemical substances	17-12
Variety (VAR).....	17-12
Primary and export grade determinants table	17-13
Corn, Canada Western/Canada Eastern Yellow, White or Mixed (CW/CE)	17-13
Export shipments	17-14
Grading.....	17-14
Cracked corn and foreign material (CCFM)	17-14

Determination of dockage

Definitions

In Eastern Canada the terms of delivery or terms of a contract will determine if dockage is to be assessed. In Western Canada dockage may only be assessed on tough, damp, moist or wet corn. The CGC will only assess dockage upon request. Dockage is assessed and recorded to the nearest 0.1%. Once dockage material has been removed it will not be reintroduced to the sample to determine the grade.

Cracked Corn and Foreign Material (CCFM) is a grading factor and will be assessed when determining the grade.

Dockage is defined under the Canada Grain Act as “any material intermixed with a parcel of grain, other than kernels of grain of a standard of quality fixed by or under this Act for a grade of that grain that must and can be separated from the parcel of grain before that grade can be assigned to the grain”. Dockage is removed by following the cleaning procedures described in this section of the manual.

The sample as it arrives is referred to as the uncleaned or dirty sample. Its weight is the **gross weight** of the sample. Dockage is assessed on the gross weight of the sample.

Dockage not reported

- ▲ **Important:** Dockage is not reported for
 - Corn, Sample CE Account Fireburnt
 - Corn, Sample Salvage
 - Corn, Sample Condemned
 - Unofficial samples declared as processed

Normal cleaning procedures:

- ▲ **Important:** Wear gloves and a mask to handle any sample which you suspect may contain hazardous substances.

Assess dockage before assessing test weight.

Procedures for normal cleaning of corn require

- No. 12 round-hole hand sieve for corn with 25% moisture or less
 - No. 14 round-hole hand sieve for corn over 25% moisture
1. Using a Boerner-type divider, divide the uncleaned sample to obtain a representative portion.
 - Official samples shall be at least 1kg.
 - Unofficial samples shall be at least 1 kg.
 2. Sieve approximately 250 g at a time on the appropriate hand sieve until all possible material has fallen through the sieve.
 3. Handpick material remaining on top of the sieve as described under *Composition of dockage*.

Composition of dockage

- All material which passes through the No. 12 or No. 14 round-hole sieve
- All foreign material and pieces of cob handpicked from the sample, excluding stones

Estimating test weight of well-matured corn after drying

Note: Test weight is determined on corn prior to removal of cracked corn and foreign material. When the Terms of Delivery or terms of a contract state that dockage can be deducted then test weight is determined after the removal of dockage.

Corn samples that contain a high percentage of moisture typically show an increase in test weight after drying. Use the following table to predict the test weight of well-matured corn before drying.

- ▲ **Important:** This is only a guide, and works only with well matured corn. Samples should be checked yearly to ensure that the formula applies for that crop condition.

Moisture range	Amount to add	
	kg/hL	g/0.5 L
15.8 - 16.4	0.5	2.6
16.5 - 16.9	1.0	5.2
17.0 - 17.3	2.0	10.5
17.4 - 17.6	2.1	11.0
17.7 - 17.9	2.2	11.5
18.0 - 18.3	2.3	12.0
18.4 - 18.6	2.4	12.5
18.7 - 18.9	2.5	13.0

1. Find the moisture range for the test weight of the sample.
2. Add the amount for that moisture range in the appropriate units, either kg/hL or g/0.5 L.

- ▲ **Important:** Corn samples are not to be downgraded on the basis of an “estimated” test weight. Questionable samples must be dried by exposure and then have the test weight determined.

For example,

The moisture for the tested sample is 17.5%, and the test weight is in kg/hL.

Moisture range	Amount to add	
	kg/hL	g/0.5 L
17.4 - 17.6	2.1	11.0

Add 2.1 to the test weight in kg/hl.

Grading

Important definitions

Net weight of sample

The sample after cleaning and removal of dockage is referred to as the cleaned sample. Its weight is the net weight of the sample. Percentages by weight for grading refer to percentages of the net weight.

Hazardous substances in samples

Wear gloves and a mask to handle any samples that you suspect may contain hazardous substances. Hazardous substances are defined in section 1 of the Canada Grain Regulations as “any pesticide, desiccant or inoculant”.

Processed sample

An unofficial sample of grain declared to be conditioned or cleaned to meet end user specifications, and whereas, the determination of dockage and/or determination of commercially clean are not performed or reported.

Rounding rules

Rounding rules are outlined in [schedule 3 of the Canada Grain Regulations](#). When official inspection results are expressed numerically, they should be expressed to the same decimal precision as the applicable tolerance in the primary and export grade determinants table.

Non-registered varieties

Where grain of any kind is not a registered variety under the Seeds Act, no person shall, except with the permission of the Canadian Grain Commission, assign a statutory grade to that grain which is higher than the lowest grade established by regulation for that kind of grain. An [order](#) is issued annually to allow non-registered varieties of corn to be graded higher than the lowest statutory grade.

Representative portions for grading

All grading is done on representative portions divided down from the clean sample, using a Boerner-type divider.

The optimum representative portion is the representative sample size within the minimum and maximum range used to obtain the most accurate result when assessing an objective factor. It is determined by taking into consideration the tolerance and concentration of the objective factor being assessed.

Representative portion of corn for grading (in grams)

Grading factor	Sample portion size range	
	Minimum	Maximum
Caramelized kernels	100 g	250 g
Classes	100 g	working sample
Cracked corn and foreign material	working sample	working sample
Damage	100 g	250 g
Excreta	working sample	working sample
Fertilizer pellets	working sample	working sample
Fireburnt	working sample	working sample
Heated and rotted	100 g	working sample
Odour	working sample	working sample
Stones	working sample	working sample
Treated seed	working sample	working sample

Grading factors

Blue-eye mould (BEM)

Germs of kernels appear dark blue with mould, or there may be just a visible mouldy blue streak under the hull of the germ. In the second case, peel back the hull from the germ to examine the germ.

Blue-eye mould is included in the tolerance for *Damage*.

Caramelized kernels

Caramelized kernels are kernels that were very immature when dried at a high temperature in a dryer, and the heat has turned the kernel to a scorched colour similar to that of heated kernels. The outer hull of the kernel may be peeled off showing a slightly damaged kernel inside. These kernels are classed as *Damaged*.

Classes

Corn is classed as yellow, white, or mixed. The class forms part of the grade name; for example, *Corn, Sample CW Yellow Account Heated*.

Samples of yellow and white corn containing less than 95% of one class are designated *Mixed*; for example, *Corn No. 1 CE Mixed*.

Contaminated grain

- ▲ **Important:** Wear gloves and a mask to handle any sample that is suspected of containing contaminated grain.

Grain is contaminated for the purposes of the *Canada Grain Act* if the grain contains any substance in sufficient quantity that the grain is either

- (a) adulterated for the purposes of the *Food and Drugs Act*; or
- (b) contaminated within the meaning of the regulations made under section 51 of the *Safe Foods for Canadians Act*.

Procedures

If a sample is suspected of being contaminated, the sample should be submitted to the Canadian Grain Commission. Determination as to whether grain is contaminated will be made by the Grain Research Laboratory in consultation with the Chief Grain Inspector for Canada. Samples deemed to be contaminated are graded: *Corn, Sample Condemned*.

Cracked corn and foreign material (CCFM)

Cracked corn and foreign material includes any of the following:

- All material including kernels and pieces of kernels of corn or any other grains which pass through a No. 12 round-hole sieve or, for samples with a moisture level over 25.0%, through the No. 14 round-hole sieve
- All foreign material other than stones handpicked from the sample, including pieces of cobs that were not removed by sieving

Procedures

▲ **Important:** Follow procedures for assessing dockage.

Note: Because breakage occurs during handling at terminal elevators, round down percentages by weight of CCFM to the nearest whole number on all officially sampled and inspected domestic or export shipments from a terminal elevator.

For example, a sample containing 4.7% CCFM by weight is recorded as containing 4% CCFM for grading purposes only on officially sampled and inspected shipments from a terminal elevator.

Damage (DMG)

Damaged kernels include whole kernels or pieces of kernels which are

- Affected by blue-eye mould and other types of moulds
 - Sprouted
 - Ground-damaged
 - Weathered
 - Diseased
 - Frosted
 - Scorched, from a drier
 - Heated, naturally, or from a drier, or caramelized
 - Rotted
-

Earth pellets (EP)

- Hard earth pellets are pellets that do not crumble under light pressure.
See *Stones*.
 - Soft earth pellets are pellets that crumble under light pressure.
See *Cracked corn and foreign material*.
-

Excreta (EXCR)

Excrement from any animal including mammals, birds and insects.

▲ **Important:** Wear gloves and a mask to handle any samples that you suspect may contain excreta.

Extraneous material

Can be defined as glass, metal, wood, plastic or any other material not already defined in the Official Grain Grading Guide.

Fertilizer pellets (FERT PLTS)

Fertilizer pellets are a manufactured plant nutrient product used by producers in the production of grain. They are typically small, round or irregular shaped and usually white, grey, brown, pink or reddish in colour.

Procedures

- Handpick any fertilizer pellets and determine the concentration basis the net working sample.
 - Fertilizer pellets are assessed as stones when the concentration does not exceed 1.0% of the net sample weight.
 - Samples containing fertilizer pellets in excess of 1.0% of the net sample weight are graded *Corn, Held IP Suspect Contaminated Grain*.
-

Fireburnt (FBNT)

Fireburnt kernels are kernels charred or scorched by fire. A cross-section of a fireburnt kernel resembles charcoal with numerous air holes. The air holes result in a low weight kernel which crumbles easily under pressure.

Procedures

Samples of corn containing fireburnt kernels are graded *Corn, Sample CW/CE (class) Account /Fireburnt*.

Foreign material (FM)

See *Cracked corn and foreign material (CCFM)*.

Heated (HTD)

Heated kernels have at least one of the following characteristics:

- Whole kernels or pieces of kernels which range in colour from amber to dark brown over the entire kernel
- Kernels which are totally discoloured by fermentation and show no natural colour on the crowns or dorsals, or both
- The germ of the kernel is amber to dark brown and is severely puffed in the germ area when heated in a drier
- Heated seeds of other grains are included in the tolerance for *Heated*

If kernels exhibit none of the above characteristics, but are not whole or sound, they are classed as *Damaged*.

Note: Cracked corn and foreign material that is heated is included with heated corn for grade assessment.

Odour (ODOR)

There is no numeric tolerance for odour. Consider

- The basic quality of the sample
- The type and degree of the odour
- The presence of visible residue causing the odour

Grains grading No. 1 through 4 must be cool and sweet. Corn grading No. 5 may have a slight odour associated with the low quality, but the odour cannot be sour or musty.

If odour is the grade determinant and there is . . .	Then the grade is . . .
An excessive objectionable odour not associated with the quality of the grain, but not heated or fireburnt	<i>Corn, Sample CW/CE Account Odour</i>
An excessive heated odour	<i>Corn, Sample CW/CE Account Heated</i>
An excessive fireburnt odour	<i>Corn, Sample CW/CE Account Fireburnt</i>

Rotted (ROT)

Rotted kernels are whole kernels or pieces of kernels which are visibly in advanced stages of decomposition and feel spongy under pressure. Rotted kernels are included in the percentage of heated kernels for grade assessment.

Stones (STNS)

Stones are hard shale, coal, hard earth pellets, and any other nontoxic materials of similar consistency. Fertilizer pellets are assessed as stones when constituting 1.0% or less of the net sample weight. (See *Fertilizer pellets* for specific procedures to be followed when samples contain fertilizer pellets.)

Procedures

1. Handpick stones from a representative portion of the cleaned sample.
2. Determine stone concentration in the net sample.
 - Samples of grain grown in western Canada samples of grain containing stones in excess of “basic grade” tolerances, up to 2.5% are graded *Corn, Rejected “basic grade” Account Stones*. The “basic grade” refers to a grade established in the Canada Grain Regulations (grades listed in the first column in grade determinant tables) that would have been assigned to the sample if it contained no stones.
 - Samples of grain grown in eastern Canada samples of grain containing stones in excess of grade tolerances are degraded to lower grades. Samples containing stones in excess of the tolerance of the lowest grade established by regulation up to 2.5% are graded *Corn, Sample CE (class) Account Stones*.
 - Samples of western and eastern Canadian grain containing more than 2.5% stones are graded *Corn, Sample Salvage*.

Examples: Western Canada

Excerpt from grade determinant tables for
Corn, Canada Western, White, Yellow or Mixed

Grade name	Stones %
No. 1 CW	0.15
No. 2 CW	0.15
No. 3 CW	0.15
No. 4 CW	0.15
No. 5 CW	0.15

Basic grade:..... *Corn, No. 2 CW Yellow*

Reason for basic grade:..... 3% Cracked corn and foreign material

If the above sample contained	Grade in Western Canada
0.5% stones	<i>Corn, Rejected No. 2 CW Yellow Account Stones</i>
3.0% stones	<i>Corn, Sample Salvage</i>

Examples: Eastern Canada

Excerpt from grade determinants table for
Corn, Canada Eastern, White, Yellow or Mixed

Grade name	Stones %
No. 1 CE	0.15
No. 2 CE	0.15
No. 3 CE	0.15
No. 4 CE	0.15
No. 5 CE	0.15

Basic grade:..... *Corn, No. 2 CE Yellow*

Reason for basic grade:..... 3% Cracked corn and foreign material

If the above sample contained	Grade in Eastern Canada
0.5% stones	<i>Corn, Sample CE Yellow Account Stones</i>
3.0% stones	<i>Corn, Sample Salvage</i>

Test weight (TWT)

Test weight is the weight of a known volume of grain expressed in kilograms per hectolitre. For procedures, see Chapter 1 of this guide, *Test weight*.

Test weight on corn is determined prior to removal of cracked corn and foreign material. When the Terms of Delivery or terms of a contract state that dockage can be deducted, then test weight is determined after the removal of dockage.

Treated seed and other chemical substances**Treated seed**

Treated seed is grain that has been adulterated with an agricultural chemical for agronomic purposes. The types of agricultural chemicals used to treat seed include pesticides, fungicides and inoculants. These seed dressings contain a dye to render the treated seed visually conspicuous. The colour of the dye varies depending upon the type of treatment and the type of grain. The current Canadian colour standard for pesticide and fungicide seed treatments for cereal (including corn) is red or pink. The colour standard for pesticide and fungicide seed treatments for canola is blue; however, green has also been used. Pulse crop (including soybeans) pesticide and fungicide seed treatments are typically blue or green. The coatings or stains may appear greasy or powdery and the surface area covered may range from tiny flecks to complete coverage.

Other chemical substances

Other chemical substances refers to any chemical residues either adhering to the kernel or remaining in the sample and to samples having a chemical odour of any kind.

- ▲ **Important:** Wear gloves and a mask to handle any samples that you suspect may contain contaminated grain.

Procedures

If a sample is suspected of being coated with a pesticide, desiccant, inoculant or if the sample contains evidence of any foreign chemical substance other than fertilizer pellets, the sample shall be graded *Corn, Held IP Suspect Contaminated Grain*.

Variety (VAR)

Corn is graded without reference to variety.

Primary and export grade determinants table

Corn, Canada Western/Canada Eastern Yellow, White or Mixed (CW/CE)

Grade name	Standard of quality			Damage		Stones %	Foreign material		Other classes %
	Minimum test weight kg/hL (g/0.5 L)	Variety	Degree of soundness	Heated %	Total %		Excreta %	Total Cracked corn and foreign material %	
No. 1 CW/CE	68 (343)	Any variety of corn	Cool and sweet, uniform size	0.1	3	0.15	0.02	2	5
No. 2 CW/CE	66 (333)	Any variety of corn	Cool and sweet	0.2	5	0.15	0.02	3	5
No. 3 CW/CE	64 (322)	Any variety of corn	Cool and sweet	0.5	7	0.15	0.02	5	5
No. 4 CW/CE	62 (311)	Any variety of corn	Cool and sweet	1.0	10	0.15	0.02	7	5
No. 5 CW/CE	58 (291)	Any variety of corn	May have a slight odour, not sour or musty	3.0	15	0.15	0.02	12	5
Grade, if No. 5 specs not met	<i>Corn, Sample CW/CE (class) Account Light Weight</i>			<i>Corn, Sample CW/CE (class) Account Heated</i>	<i>Corn, Sample CW/CE (class) Account Damaged</i>	<i>2.5% or less—Corn, Rejected (grade) (class) Account Stones or Corn, Sample CE (class) Account Stones Over 2.5%—Corn, Sample Salvage</i>	<i>Corn, Sample CW/CE (class) Account Excreta</i>	<i>50% or less – Corn Sample CW/CE (class) Account CCFM Over 50% - sample Cracked Corn and Foreign Material</i>	<i>Over 5%—Use all other grading criteria and grade as Corn (grade) Mixed</i>

Note: The colour is added to the grade name.

Export shipments

Grading

Corn on export is graded in accordance with primary grade standards and specifications.

Cracked corn and foreign material (CCFM)

Because breakage occurs during handling at terminal elevators, round down percentages by weight of CCFM to the nearest whole number on all officially sampled and inspected domestic or export shipments from a terminal elevator.

For example, a sample containing 4.7% CCFM by weight is recorded as containing 4% CCFM for grading purposes only on officially sampled and inspected shipments from a terminal elevator.